

MONOMIAL MULTIPLE STRUCTURES

JON EIVIND VATNE

ABSTRACT. In this paper we study monomial multiple structures on a linear subspace of codimension two in projective space. We show that these structures determine smooth points in their respective Hilbert schemes, with (smooth) neighbourhoods of two such points intersecting if their Hilbert functions are equal. We generalize a construction for multiple structures on points in the plane to this setting, giving a kind of product of monomial multiple structures. For points, this construction can be found in Nakajima's book [5]. The tools we use for studying multiple structures are developed in [6] and [7].

1. INTRODUCTION

The monomial multiple structures are among the best understood multiple structures. They can be visualized using Young diagrams, their invariants are easily calculated, and their “inductive construction” (see below) is quite simple. In this paper we focus on monomial multiple structures on linear subspaces of codimension two in projective space satisfying the Cohen-Macaulay property. Cohen-Macaulay always means *locally* Cohen-Macaulay.

Let us first summarize the method for constructing multiple structures from [6] and [7], adapted to our special situation: Let $X = \mathbb{P}^n \subset \mathbb{P}^{n+m}$ be a linear subspace, $X^{(i)} \subset \mathbb{P}^{n+m}$ the i 'th infinitesimal neighbourhood of X , and Y a Cohen-Macaulay multiple structure with $Y_{\text{red}} = X$, whose ideal is generated by monomials. Then there is a filtration of Y

$$X = Y_0 \subset Y_1 = Y \cap X^{(1)} \subset \cdots \subset Y_{k-1} = Y \cap X^{(k-1)} \subset Y_k = Y \cap X^{(k)} = Y$$

for some k . Since Y is a monomial Cohen-Macaulay structure, the same is true for each term in this filtration, but in general the terms will not have such good properties. To keep the terminology the same as in [6] and [7] we will therefore refer to this filtration as the S_1 -filtration of Y .

Let $\mathcal{I}_i$ be the Ideal of Y_i . Then there are two short exact sequences

$$(1) \quad 0 \rightarrow \mathcal{I}_{j+1}/\mathcal{I}_X \mathcal{I}_j \rightarrow \mathcal{I}_j/\mathcal{I}_X \mathcal{I}_j \rightarrow \mathcal{L}_j \rightarrow 0$$

and

$$(2) \quad 0 \rightarrow \mathcal{L}_j \rightarrow \mathcal{O}_{Y_{j+1}} \rightarrow \mathcal{O}_{Y_j} \rightarrow 0.$$

Here the $\mathcal{O}_X$ -Module $\mathcal{L}_j$ is defined by the first exact sequence, and the $\mathcal{O}_{Y_i}$ are $\mathcal{O}_X$ -Modules by restricting a projection onto X to each Y_i . The Hilbert polynomial of Y can be calculated as

$$\text{Hilb}(Y) = \text{Hilb}(X) + \sum_j \text{Hilb}(\mathcal{L}_j).$$

In general, the terms $\mathcal{L}_j$ will be torsion free, but in the monomial case they are actually *locally free*.


The paper is organized as follows: In Section 2 we give the basic description of monomial multiple structures in terms of Young diagrams. In Section 3 we show that the points in the Hilbert scheme determined by monomial multiple structures in codimension two are smooth, and relate neighbourhoods of these points for different multiple structures. In Section 4 we describe a procedure for taking a kind of product of two monomial multiple structures in codimension two, corresponding to a natural operation on Young diagrams.

Acknowledgements This article is based on parts of my Doctorate thesis [6]. I would like to thank the group in algebraic geometry at the University of Bergen, G.Fløystad, T.Johnsen, A.L.Knutsen, S.-A.Strømme, J.-M.Økland and most of all my advisor A.Holme. I will also express my gratitude to the evaluation committee for many useful comments and suggestions: T.Johnsen, K.Ranestad and F.-O.Schreyer.


2. YOUNG DIAGRAMS AND MONOMIAL IDEALS

Definition 2.1. An n -dimensional *Young diagram* is a finite array of unit boxes in the first hyperquadrant in Euclidean n -space, with corners in the integer grid, such that wherever there is a box, there is always a box in the place immediately below it in each direction. When we speak of two-dimensional Young diagrams we will suppress the number “two” and speak simply of Young diagrams, and we will use terms like “up, down, left and right”. The *weight* of a box is the sum of the coordinates of the innermost corner in the diagram.

It is well known that Young diagrams correspond to partitions: The parts of the partition are the number of boxes in the row corresponding to the place of the part in the partition, where the partition is ordered by (weakly) decreasing magnitude. As an example, to the partition $(4, 4, 3, 1)$ of 12 corresponds the Young diagram



We number boxes as follows: the box $(r_1, \dots, r_m)$ of a Young diagram is the box with this integer point as its lower corner (lower in all directions). For instance,




Young diagrams arise from Cohen-Macaulay monomial ideals whose radical is the ideal of a linear subspace. We will constantly use the following elementary lemma without notice:

Lemma 2.2. *Let $I_X = (x_1, \dots, x_m) \subset k[x_1, \dots, x_m, z_0, \dots, z_n]$ be the (saturated) ideal of a linear subspace $X = \mathbb{P}^n \subset \mathbb{P}^{n+m}$. Let $J = I_Y$ be a (saturated) monomial ideal whose radical is I , i.e. $Y_{\text{red}} = X$. Then the following are equivalent:*

- Y satisfies (S_1) .
- Y is (locally) Cohen-Macaulay.
- Y is arithmetically Cohen-Macaulay.


Proof. It is obviously enough to show that Y is arithmetically Cohen-Macaulay if Y satisfies (S_1) . So assume this. Now each member of a minimal monomial generator set of J looks like $x_1^{a_1} \cdots x_m^{a_m} z_0^{b_0} \cdots z_n^{b_n}$ for some non-negative integers a_i and b_j . If any b_j is non-zero z_j is necessarily contained in an associated prime ideal of Y , but by assumption Y does not have any embedded primes. Thus all $b_j = 0$ and $z_0, \dots, z_n$ is a *global* regular sequence on Y and Y is arithmetically Cohen-Macaulay. $\square$

For the construction of associated Young diagrams, we start with $X = \mathbb{P}^n \subset \mathbb{P}^{n+2}$, with ideal $I_X = (x, y) \subset S := k[x, y, z_0, \dots, z_n]$, and we consider (Cohen-Macaulay) monomial ideals $J \subset S$ whose radical is I_X . To such an ideal J we associate a Young diagram. First of all we will think of x, y as being directions or axes in the Euclidean plane where we draw our Young diagrams, x pointing to the right and y pointing up, as usual. Then each monomial in x, y gives an integer point in the first quadrant, $x^r y^s \leftrightarrow (r, s)$. We write the monomial in that position:



Now, consider the monomial ideal J . Since it is an ideal, whenever a monomial $x^r y^s$ lies in J , all multiples $x^{r+u} y^{s+v}$ also lie in J . Saying that the radical of J is (x, y) is the same as saying that there are only finitely many pairs (r, s) such that

$x^r y^s \notin J$. Put together, the monomials not in J form a Young diagram. For instance, $J = (x^4, x^3 y^2, x^2 y^3, y^4)$ gives the Young diagram (where we write the generators of the ideal in their respective positions outside the Young diagram)



Obviously this is a bijective correspondence between Cohen-Macaulay monomial ideals in two variables and Young diagrams. Similarly,

Proposition 2.3. *There is a bijective correspondence between Cohen-Macaulay monomial ideals in l variables having a given linear subspace of codimension l as support, and l -dimensional Young diagrams. Under this correspondence, the number of boxes in the Young diagram is the multiplicity of the scheme defined by the corresponding ideal. The correspondence is inclusion-reversing.*

Proof. The correspondence is as above. The equality between the number of variables occurring in the ideal and the codimension ensures that there are only finitely many boxes.

Each box in the Young diagram gives a monomial not in J . These monomials form a basis for the homogeneous coordinate ring S/J as an S/I_X -module by Theorem 3.3. The number of basis elements is equal to the rank of this module, or equivalently, the rank of the structure sheaf as a Module over the structure sheaf of the reduced subscheme. This rank is the multiplicity.

If one Young diagram contains another, the corresponding ideal has fewer monomials than the second one's, and vice versa. $\square$

The statement about multiplicities in the proposition will be sharpened in the following sections; we will see, that the entire Hilbert polynomial and Hilbert function can be read off the diagram.

Some standard operations on ideals are easily described using the associated Young diagrams:


Proposition 2.4. *Let I, J be Cohen-Macaulay monomial ideals in $x_1, \dots, x_m$ with radical $(x_1, \dots, x_m)$, with associated Young diagrams T and S .*

- *The Young diagram of $I + J$ is $T \cap S$.*
- *The Young diagram of $I \cap J$ is $T \cup S$.*
- *The Young diagram of the ideal of the k th infinitesimal neighbourhood is given by all boxes on or below the k th diagonal.*
- *The Young diagram of the k th part of the S_1 -filtration of I , I_k , is given by the part of the Young diagram of I on or below the k th diagonal.*


By “the k th diagonal” we mean the set of boxes $r_1, \dots, r_m$ such that the weight $r_1 + \dots + r_m$ is constant, equal to k . In dimension two a diagonal is thus an ordinary diagonal going from the upper left to the lower right. In higher dimensions it is a “large diagonal”.

Proof. A monomial not in $I + J$ is a monomial neither in I nor in J , so it corresponds to a box in both T and S . Similarly, a monomial in $I \cap J$ is in both I and J , so one not in $I \cap J$ is outside I or outside J . Thus the corresponding box is in either T or S . The third part follows by noting that the monomials of a given degree have the same “distance” from the origin, they have the same *weight*, so they form a diagonal. Thus the Young diagram of an infinitesimal neighbourhood is given by all boxes on or below a diagonal. The last statement is immediate from the first and the third. $\square$


Example 2.5. Let $I = (x^2, xy^2, y^3)$ and $J = (x^4, xy, y^2)$, so that $I + J = (x^2, xy, y^2)$ and $I \cap J = (x^4, x^2y, xy^2, y^3)$. First we will consider $I + J$:



Then we consider $I \cap J$:



At last we consider the S_1 filtration of J :



Proposition 2.6. *Given a Cohen-Macaulay monomial ideal I in $x_1, \dots, x_m$ with radical $(x_1, \dots, x_m)$, consider a simple thickening*

$$0 \rightarrow \mathcal{J}/\mathcal{I}\mathcal{I}_X \rightarrow \mathcal{I}/\mathcal{I}\mathcal{I}_X \rightarrow \mathcal{L} \rightarrow 0$$

where the map to $\mathcal{L} = \bigoplus \mathcal{O}_X(-a_i)$ is given by projection onto a summand generated by some of the minimal monomial generators of I . The (m -dimensional) Young diagram of $\mathcal{I}\mathcal{I}_X$ is constructed from the Young diagram of I by introducing new boxes in the inner corners. The Young diagram of the ideal J defined by the short exact sequence is constructed by introducing the boxes corresponding to the monomial generators of $\mathcal{L}$.

By inner corner of the Young diagram we mean a box not in the diagram, but such that the box immediately below it in each direction is in the diagram. There are also outer corners; those are the boxes such that the box immediately below it in each direction is outside the Young diagram, whereas the box meeting it at its corner closest to the origin is in the diagram.

Proof. We have already used the (obvious) fact that an ideal is generated by the monomials corresponding to inner corners of the diagram. Multiplying a generator with one of the generators of the ideal I_X corresponds to moving a box one step up in the direction determined by that generator. Thus a monomial not in II_X is a monomial that corresponds to a box at most one step up from a box in the diagram of I in all directions. Thus the monomials not in II_X correspond to monomials in the diagram of I or in the inner corners of that diagram. The ideal J contains the ideal II_X , but also the generators not sent to $\mathcal{L}$. Thus J corresponds to the diagram given by introducing boxes in the indicated subset of the inner corners. $\square$

3. POINTS IN HILBERT SCHEMES COMING FROM YOUNG DIAGRAMS

Proposition 3.1. *Given a Cohen-Macaulay monomial ideal I with support a linear subspace of codimension two, and with (two-dimensional) Young diagram T . Then the syzygies of I correspond to the outer corners of T . More precisely: if I has inner corners of weight n_{1j} and outer corners of weight n_{2i} , then the minimal resolution of I has the form*

$$0 \rightarrow \bigoplus_i S(-n_{2i}) \rightarrow \bigoplus_j S(-n_{1j}) \rightarrow I \rightarrow 0.$$

Note that there is one outer corner less than there are inner corners, so the ranks add up correctly.

Proof. For each outer corner, there is an inner corner to its left and another downwards. Thus there is a relation of degree equal to the weight of the outer corner between the generators corresponding to these two boxes, saying that a power of x times the monomial corresponding to the inner corner to the left is equal to a power of y times the monomial corresponding to the inner corner downwards. It is easily verified that these relations generate the full module of syzygies. $\square$

We will need these two sets of integers later on. We silently assume that they are written in decreasing order, so that it makes sense to talk about them as *sequences*.

Corollary 3.2. *The dimension of the Hilbert scheme of closed subschemes in $\mathbb{P}^{n+2}$ in the point corresponding to I is*

$$\begin{aligned} & \sum_{n_{2i} \geq n_{1j}} \binom{n_{2i} - n_{1j} + n + 2}{n + 2} + \sum_{n_{1j} \geq n_{2i}} \binom{n_{1j} - n_{2i} + n + 2}{n + 2} \\ & - \sum_{n_{2i} \geq n_{2j}} \binom{n_{2i} - n_{2j} + n + 2}{n + 2} - \sum_{n_{1j} \geq n_{1i}} \binom{n_{1j} - n_{1i} + n + 2}{n + 2} + 1. \end{aligned}$$

The Hilbert scheme is smooth in the point determined by I .

Proof. The displayed equation is from Ellingsrud, [3], where the condition is that the ideal is of codimension two and is arithmetically Cohen-Macaulay. This holds in our situation because of Lemma 2.2. $\square$

Theorem 3.3. *Given an m -dimensional Young diagram with corresponding multiple scheme Y with support X , the structure sheaf of Y , considered as an $\mathcal{O}_X$ -Module, is*

$$\mathcal{O}_Y \cong \bigoplus_B \mathcal{O}_X(-w_B).$$

Here the sum is over the boxes B in the diagram, and w_B is the weight of the box B . Introduce multiplicative structure on $\bigoplus_B \mathcal{O}_X(-w_B)$ by introducing the monomial corresponding to B as generator for $\mathcal{O}_X(-w_B)$. Then the above isomorphism becomes an isomorphism of $\mathcal{O}_X$ -Algebras.

Proof. The proof is by examining what happens with the Young diagram when we perform our standard thickenings of multiple structures. So let I be a monomial ideal, $I_X = (x_1, \dots, x_m)$ the ideal of the support with $\mathcal{I}, \mathcal{I}_X$ the associated sheaves. Then our standard construction gives us

$$\bigoplus_{j=1}^s \mathcal{O}_X(-a_j) \cong \mathcal{I}/\mathcal{I}I_X$$

where the $a_j = w_B$ are degrees of the generators of $\mathcal{I}$, and are thus given by sums $r_1 + \dots + r_m$ for generating monomials $x_1^{r_1} \dots x_m^{r_m}$. Since the generating monomials are represented by the corners of the Young diagram, we see that in the standard exact sequence

$$0 \rightarrow \bigoplus_{j=t+1}^s \mathcal{O}_X(-a_j) \rightarrow \mathcal{I}/\mathcal{I}I_X \rightarrow \bigoplus_{j=1}^t \mathcal{O}_X(-a_j) \rightarrow 0$$

the ideal J with $\mathcal{J}/\mathcal{I}I_X = \bigoplus_{j=t+1}^s \mathcal{O}_X(-a_j)$ has Young diagram obtained from the Young diagram of I by adding the boxes corresponding to $\bigoplus_{j=1}^t \mathcal{O}_X(-a_j)$. Now this gives an additional summand $\bigoplus \mathcal{O}_X(-a_j)$ to the structure sheaf by the short exact sequence 2 (which splits in this our case). Using this as an induction step, the theorem is proven (the case corresponding to a single box being trivial).

The multiplicative structure is easily seen to be the same on both sides. $\square$

Remark 3.4. The multiplicative structure on the sum $\bigoplus_B \mathcal{O}_X(-w_B)$ is a multiplicative structure on the sum of the $\mathcal{L}_j$ s of the S_1 -filtration. A similar (but more general) construction was considered for curves by Bănică and Forster, [1] and [2], and also by Manolache in higher dimensions, see [4].

Corollary 3.5. *The Hilbert polynomial of the scheme associated to an m -dimensional Young diagram can be computed as follows: Define*

$$b_i(d) = \binom{n+d-i}{n} = \chi(\mathcal{O}_{\mathbb{P}^n}(d-i)).$$

Then the Hilbert polynomial is

$$\sum_B b_{w_B}$$

the sum over all boxes B in the diagram.

Remark 3.6. Another way of stating this corollary is to say that the Hilbert polynomial of the multiple structure is equal to the Hilbert polynomial of the sum of the invertible sheaves $\mathcal{O}_X(-w_B)$ from the theorem. In this formulation, the statement can be strengthened to an equality of Hilbert *functions*.

Proposition 3.7. *All the Cohen-Macaulay monomial ideals with a given codimension two linear subvariety as support, with a given Hilbert function, lie in the same irreducible component of the Hilbert scheme.*

Example 3.8. We cannot replace “Hilbert function” by “Hilbert polynomial” in the previous proposition. Consider for example the two ideals $I = (x^5, x^4y, y^2)$ and $I' = (x^6, x^2y, xy^2, y^3)$ in $k[x, y, z, w]$. The two multiple schemes defined on the line $x = y = 0$ by these ideals have the same Hilbert polynomials, but different Hilbert functions. Using Corollary 3.2 we find that the dimension of the Hilbert scheme in the point corresponding to I is 38, whereas it is 39 in the point corresponding to I' .

In order to prove this proposition, we need some combinatorial definitions, as well as another result from [3].

Definition 3.9. Define the equivalence relation R on the set of Young diagrams by TRT' if and only if the ideals defined by T and T' have the same Hilbert function. By Theorem 3.5 and Remark 3.6, this is equivalent to saying that T and T' have the same number of boxes in each diagonal.

Remark 3.10. For Young diagrams that are “small” with respect to the dimension, this is also equivalent to having the same Hilbert polynomial. In fact, given two monomial ideals I, I' with the same Hilbert polynomial, but with different Hilbert functions, we can regard them as ideals in a polynomial ring with more variables (i.e. we can consider their projective cones). Then, for a sufficiently high number of variables, the Hilbert polynomials will also be different. Basically, this is because there are relations between the classes of sums of line bundles in the Grothendieck group of $\mathbb{P}^N$, but these relations cannot be extended indefinitely. Thus Young diagrams differing by boxes corresponding to line bundles satisfying some identities in low dimensions will not be determined by their Hilbert polynomial in these low dimensions.

Definition 3.11. Define the equivalence relation r on the set of Young diagrams as follows: By Proposition 3.1 each Young diagram determines the resolution

$$0 \rightarrow \bigoplus_{i \in A} S(-n_{2i}) \rightarrow \bigoplus_{j \in B} S(-n_{1j}) \rightarrow I \rightarrow 0$$

for some index sets A and B . Consider the two sequences of integers $(\{n_{2i}\}, \{n_{1j}\})$. We say that two such pairs $(\{n_{2i}\}, \{n_{1j}\})$ and $(\{n'_{2i'}\}, \{n'_{1j'}\})$ are primitively equivalent if there exists subsets $A_0 \subset A, B_0 \subset B$ and a bijection $\sigma : A_0 \rightarrow B_0$ such that

$n_{2i} = n_{1\sigma(i)}$ holds for all $i \in A_0$, and $\{n'_{2i'}\} = \{n_{2i}\} \setminus A_0$, $\{n'_{1j'}\} = \{n_{1j}\} \setminus B_0$. We say that TrT' if the sets of integers determined by T and T' are equivalent by the equivalence relation generated by the primitive equivalences above. This should be compared with Ellingsrud's "prolongement", see [3], page 424, on which it is based.

What pairs of sequences of integers can appear?


Lemma 3.12. a) *A pair of sequences of integers $(\{a_i\}, \{b_j\})$ is associated to some Young diagram if and only if*

- i) *There is exactly one more a_i then there are b_j s.*
- ii) *$b_i > a_i \geq a_{i+1}$ for all i .*
- iii) *$\sum_i a_i = \sum_j b_j$.*

b) *Any Young diagram is r -equivalent to a Young diagram without any equalities of the kind $a_i = b_j$.*

In part a)ii) the inequality $a_i \geq a_{i+1}$, a part of our assumptions, is included only for emphasis.

Proof. For part a), the necessity of the conditions is obvious. Conversely, given such a pair we can construct the Young diagram (where we write the weights on inner and outer corners)



The box corresponding to the number a_1 has coordinates $(0, a_1)$. The box corresponding to the number b_1 has coordinates $(b_1 - a_1, a_1)$. The box corresponding to the number a_2 has coordinates $(b_1 - a_1, a_1 - b_1 + a_2)$. Continuing in this way, we finally come to the box corresponding to the number a_s , which has coordinates

$$(b_1 - a_1 + b_2 - a_2 + \cdots + b_{s-1} - a_{s-1}, a_1 - b_1 + a_2 - b_2 + \cdots + a_s) = (a_s, 0)$$

The conditions ensure that the first coordinates of the a_i (equal to the first coordinates of the b_{i-1}) form an increasing sequence from 0 to a_s , whereas the second coordinates of the a_i (equal to the second coordinates of the b_i) form a decreasing sequence from a_1 to 0.

Note that the Young diagram constructed in this way is only one of many different Young diagrams determining these sequences.

Part b) follows from part a): just remove any pairs $a_i = b_j$ and write the resulting sequences in order of decreasing magnitude. The conditions i) – iii) are still verified. $\square$

Now the result from [3], translated to the (special) case of monomial ideals:

Proposition 3.13. *Given Young diagrams T and T' . Let P, P' be the corresponding points in the Hilbert scheme. Then there are smooth open neighbourhoods around P and P' that intersect if and only if TrT' . For a given Hilbert function, the union of all open sets thus determined for all Young diagrams is a smooth, connected scheme.*

Proof. See [3] page 426.(Theorem 2.iii). Note that we only use a small part of this theorem.

In the statement of the theorem of Ellingsrud's, in order to get a non-empty intersection, we need to know that there is a scheme with the common refinement as the pair of sequences arising from its minimal resolution. This is automatic in the case of a given Hilbert function, since (by Lemma 3.12) there is a monomial ideal with the same Hilbert function, without any equalities like $n_{2i} = n_{1j}$. $\square$

Proposition 3.14. *The equivalence relations r and R are the same.*


Proof. Since the Hilbert function of I can be computed from the Young diagram by Remark 3.6, it is easily seen that $TrT' \Rightarrow TRT'$.

For the other implication, note that by Remark 3.10, if the Hilbert functions of two monomial ideals are different, then the zero schemes are (successive) hyperplane sections of monomial schemes with different Hilbert polynomials. These will nevertheless have the same Young diagrams as the original monomial ideals. It easily follows that these Young diagrams are different. Thus $TRT' \Rightarrow TrT'$. $\square$

With these propositions, the proof of Proposition 3.7 is complete.

4. SUMS OF YOUNG DIAGRAMS, PRODUCTS OF MULTIPLE STRUCTURES

There is a simple operation on two-dimensional Young diagrams that we are going to need, see Chapter 7 of Nakajima's book [5]. Given two partitions $\lambda = (\lambda_0 \geq \lambda_1 \geq \dots \geq \lambda_k \geq 0)$ and $\mu = (\mu_0 \geq \mu_1 \geq \dots \geq \mu_k \geq 0)$ (we extend the shortest partition by adding zeroes, if necessary, so that they have the same length), we can form their partwise sum $(\lambda_1 + \mu_1 \geq \dots \geq \lambda_k + \mu_k)$. The result is again a partition. The operation on Young diagrams is to add the number of boxes in each row. For instance $(4, 4, 3, 2) + (3, 3, 1) = (4, 4, 3, 2) + (3, 3, 1, 0) = (7, 7, 4, 2)$ or



This operation has several good properties:

Proposition 4.1. *The operation described above is associative, commutative and has unit given by the empty diagram. Thus the set of all Young diagrams becomes an abelian monoid. This monoid is generated by Young diagrams with only one column.*

Proof. The fact that we get an abelian monoid is trivial. The last statement follows since any Young diagram is the sum of its columns (or any partition is the sum of partitions of the form $(1, 1, \dots, 1)$). $\square$


Based on this operation, we will also construct a three-dimensional Young diagram as follows:

Construction 4.2. *Given two partitions λ and μ , we make a three-dimensional Young diagram with boxes*


$$\{(i, j, k) | (i, j) \in \text{the diagram of } \mu, k < \lambda_j\}$$

Remark 4.3. This is *not* a commutative construction. It depends on the order we write the two diagrams in.

This can be visualized diagrammatically: consider the skew diagram given by the right hand side minus the right member of the left side in the defining sum of two Young diagrams (where λ and μ are added in that order):



(this corresponds to the sum of $\lambda = (4, 4, 3, 2)$ and $\mu = (3, 3, 1)$ as above). The first part consists of those X s with at least one empty box to their right (a *part* is a subset of the three dimensional diagram given by all boxes with a given third coordinate). This condition means that $0 < \lambda_j$. The second part consists of those X s that have at least two empty boxes to their right, meaning $1 < \lambda_j$, and so on. Here the parts are counted from the first part with third coordinate 0, to the second part with third coordinate 1 and so on. Thus the two-dimensional parts of the three-dimensional Young diagram defined by the above equation are



In picture number i , the third space coordinate is i . Consider for example the X in position $(1, 1)$. Then $(1, 1, k)$ is in the three-dimensional diagram if and only if $k < \lambda_1 = 4$, and 4 is also the number of empty boxes to the right of this X .

Remark 4.4. If we add the two partitions in the opposite order, and carry through this construction, the result will be the mirror image of the first diagram; see Corollary 4.10.

Remark 4.5. This construction will always give a three-dimensional Young diagram. On the other hand, if the skew diagram is not made from two Young diagrams

as above, there is no reason why the result should be a three-dimensional Young diagram.

Remark 4.6. If the two partitions we add are $\lambda = (\lambda_0 \geq \lambda_1 \geq \cdots \geq \lambda_k \geq 0)$ and $\mu = (\mu_0 \geq \mu_1 \geq \cdots \geq \mu_k \geq 0)$, then the total number of boxes in the three-dimensional diagram is $\sum_l \lambda_l \mu_l$ since each box in the diagram corresponding to μ in the l th row is repeated λ_l times.

Now we consider the operation on monomial ideals corresponding to this operation on Young diagrams. For points in the plane, this was considered by Nakajima, see [5] Chapter 7. We give an example from this theory:

Example 4.7. We want to consider two points lying on the line $y = 0$ in the projective plane $\mathbb{P}^2 = \text{Proj } k[x, y, z]$. Removing the line at infinity $z = 0$, we can coordinatize this line with the remaining coordinate x , and let x_1, x_2 be the points in question. Let there be given two multiple structures on these points, with ideals $J_1 = ((x - x_1)^4, (x - x_1)^3 y^2, (x - x_1)^2 y^3, y^4)$ and $J_2 = ((x - x_2)^3, (x - x_2) y^2, y^3)$. Then the ideal of the union of the schemes is given by

$$(y^4, y^3(x - x_1)^2, y^2(x - x_1)^3(x - x_2), (x - x_1)^4(x - x_2)^3)$$

and the ideal of the special fiber, as x_2 tends to x_1 , becomes (after the simplification $x_1 = 0$)

$$(y^4, y^3 x^2, y^2 x^4, x^7).$$

The sum of the Young diagrams of the two original ideals is equal to the Young diagram of the special fibre:

$$\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} = \begin{array}{|c|c|c|c|c|c|c|} \hline \square & \square & \square & \square & & & & \\ \hline \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \square & \square & \square & \square & \square & \square & \square & \square \\ \hline \end{array}$$

The flatness of the deformation can be deduced directly from the Young diagrams: the Hilbert polynomial of the multiple point is equal to the number of boxes in the diagram, and this is “additive”.

Note that the operation is not canonical in the sense that we have chosen coordinates in the plane; the multiple structures are required to be “monomial in the coordinate y and the direction x ”.

Now it is time to describe the changes that must be made if we want to extend the results quoted above to higher dimensions. It turns out that there are very few changes, although the arguments become harder. We let $X = \mathbb{P}^n \subset \mathbb{P}^{n+2}$.

Proposition 4.8. *Consider $X = \mathbb{P}^n \subset \mathbb{P}^{n+2}$ as contained in a fixed hyperplane $H = \mathbb{P}^{n+1} \subset \mathbb{P}^{n+2}$. The ideals are $I_X = (x, y)$ and $I_H = (y)$. Let $Z = \mathbb{P}^n$ be another linear subspace, also contained in H , with ideal $I_Z = (y, z)$. Then, given Cohen-Macaulay monomial ideals in $(x, y), (z, y)$ (with these ideals as radicals), let $z \rightarrow x$. The family given by the union of the two multiple schemes on the two linear subspaces*

X and Z_t , where Z_t is determined by Z under the substitution $z \mapsto tz + (1 - t)x$, is a flat family, and the special fiber ($t = 0$) is given by adding the Young diagrams.

Proof. We can repeat the calculation from Example 4.7, and get the family we need. The difficulty lies in showing that this family is flat. For points, this was simple: the Hilbert polynomial just counts the number of boxes in the Young diagram. In this more general case the Hilbert polynomial is given by Corollary 3.5, and it is obvious from that description that when we sum Young diagrams, we don't get the sum of the Hilbert polynomials. But the difference is easily calculated.

Claim 1. Let $T' + T'' = T$ be a sum of Young diagrams. Then the difference between the Hilbert polynomial of the multiple structure corresponding to T and the sum of the Hilbert polynomials of the multiple structures corresponding to T' and T'' is given by the Hilbert polynomial of the multiple structure corresponding to the three-dimensional Young diagram from Construction 4.2.


We also need to give a geometric description of the three-dimensional Young diagram:

Claim 2. The three-dimensional Young diagram is the Young diagram of the intersection of the two multiple structures under consideration.

Given these two claims, the proposition follows from the general equality


$$\text{Hilb}(V \cup W) = \text{Hilb}(V) + \text{Hilb}(W) - \text{Hilb}(V \cap W).$$

Proof of Claim 1. Consider the following diagram, associated to the addition $T' + T'' = T$:



Here the X s in T correspond to the boxes in T'' . For each X on the right hand side we get a contribution to the Hilbert polynomial that is also found on the left hand side in the second member. Thus the difference in Hilbert polynomials must be found using the remaining boxes on the right hand side and the first member on the left hand side.

Consider any other box, and the corresponding box in T' .



The difference in Hilbert polynomials is $b_i - b_j$ where the box on the left has weight i , the box on the right has weight j . Note that $j - i = \text{number of } X\text{s in that row}$. Now there is a fundamental relation of binomial coefficients

$$b_i - b_j = \binom{n+d-i}{n} - \binom{n+d-j}{n} = \sum_{k=j}^{i-1} \binom{n-1+d-k}{n-1}.$$

For each X in the row, the marked box determines a box in the three-dimensional diagram above X , with third coordinate equal to the difference between the weight of the marked box and the weight of the rightmost box with an X . For each k between j and $i - 1$ this gives a box in the three-dimensional diagram with weight k . This explains the terms of the binomial relation. For the example marked above, we have a box in position $(4, 1)$ on the right, and $(1, 1)$ on the left. There are three X s in this row, and the three boxes in the three-dimensional diagram coming from this marked box have coordinates $(0, 1, 1)$, $(1, 1, 1)$ and $(2, 1, 1)$.

The claim follows by letting the marked box run through the skew diagram (the boxes without X s).

Proof of claim 2. A box (i, j, k) , with corresponding monomial $x^i y^j z^k$, is in the three-dimensional diagram corresponding to the sum of λ and μ if (i, j) is in the diagram μ (meaning that the monomial $x^i y^j$ is not in the ideal corresponding to μ) and if $k < \lambda_j$ (meaning that $y^j z^k$ is not in the ideal corresponding to λ .) This proves the claim. $\square$

Corollary 4.9. *The Young diagram of the intersection of two Cohen-Macaulay monomial multiple structures whose supports are linear subspaces of codimension two, contained in a common hyperplane, is given by the construction of a three-dimensional Young diagram from a pair of two-dimensional Young diagrams.*

Corollary 4.10. *The operation of constructing a three-dimensional Young diagram from a pair of two-dimensional Young diagrams does not depend on the order of the two, up to permutation of the axes.*

Remark 4.11. The permutation of the axes mentioned in the second corollary is given by permutation of the summands when adding the two Young diagrams. In the first corollary, this has the same effect; we will get the two possible choices of three-dimensional Young diagram by deciding what order we will write the intersection in.

REFERENCES

- [1] C. Bănică and O. Forster. Sur les structures multiples. Handwritten manuscript, 1981.
- [2] C. Bănică and O. Forster. Multiplicity structures on space curves. *Contemp. Math.*, 58:47–64, 1986.
- [3] G. Ellingsrud. Sur le schéma de Hilbert des variétés de codimension 2 dans $\mathbb{P}^e$ à cône de Cohen-Macaulay. *Ann.Sci. Ecole Norm. Sup.*, 8:423–432, 1975.
- [4] N. Manolache. Multiple structures on smooth support. *Math. Nachr.*, 167:157–202, 1994.

- [5] H. Nakajima. *Lectures on Hilbert Schemes of Points on Surfaces*, volume 18 of *Univ. Lecture Ser.* AMS, 1999.
- [6] J.E. Vatne. *Towards a Classification of Multiple Structures*. PhD thesis, University of Bergen, 2001. Available from <http://www.mi.uib.no/~jonev>.
- [7] J.E. Vatne. Multiple structures. preprint, math.AG/0210042, 2002.

MATEMATISK INSTITUTT, JOHS. BRUNSGT. 12, N-5008 BERGEN, NORWAY
E-mail address: `jonev@mi.uib.no`